



Smoothing effects for a transport diffusion equation

Samira Alamin Sulaiman

&

Jameelah Ebreed

Department of mathematics, Faculty of sciences, University of Zawia, Zawia
/ Libya

samira.sulaiman@zu.edu.ly

j.ebreed@zu.edu.ly



Abstract:

In this paper, we prove a smoothing effects for a transport diffusion model, with a fractional dissipation, and in the Besov spaces $B_{2,1}^0$ given in Definition 2.2.4. We will use the technic of [12].

المخلص:

في هذا البحث سوف ندرس منظومة معادلة النقل التوزيعي لمائع غير قابل للانضغاط في فضاء بزوف $B_{2,1}^0$ الدالي (تعريف 2.2.4). سوف نستعمل للبرهان الفكرة التي اعتمد عليها المؤلف الأول [12] من هذا البحث.

Keywords: Transport diffusion equation, Incompressible fluid flow, smoothing effects.

1 Introduction

In this paper, we are concerned with the initial value problem of the 2D dissipative quasi-geostrophic model

$$(1.1) \quad \begin{cases} \partial_t \theta + v \cdot \nabla \theta + (|D|^\alpha + I)\theta = f \\ \operatorname{div} v = 0, \\ \theta|_{t=0} = \theta_0, \end{cases}$$

where θ is the scalar function represents the potential temperature and the parameter $\alpha \in [0,1]$. The 2D velocity field $v = v(x, t)$, $x \in \mathbb{R}^2$, $t \in \mathbb{R}_+$, is determined by Riesz transform $R_i, \forall i = 1,2$ of θ , that is

$$v = \left(-\frac{\partial_2}{|D|} \theta, \frac{\partial_1}{|D|} \theta \right) := (-R_2 \theta, R_1 \theta).$$

The differential operator $v \cdot \nabla$ is defined respectively by

$$v \cdot \nabla = \sum_{i=1}^d v_i \partial_i.$$

The fractional differential operator $|D| = (-\Delta)^{\frac{1}{2}}$ is defined by its Fourier transform

$$\mathcal{F}(|D|u) = |\xi| \mathcal{F}(u),$$

and the operator $div v$ is defined by

$$div v = \sum_{i=1}^2 \partial_i v^i.$$

The first equation of (1) serves as a 2D models arising in geophysical fluid dynamic [11] and the second equation $div v = 0$, describe the incompressibility of the fluid.

This equation has been intensively investigated and much attention is carried to the problem of global well-posedness. For the sub-critical case ($\alpha > 1$) the theory seems to be in a satisfactory state. Indeed, global existence and uniqueness for arbitrary initial data are established in various function spaces (see for example [6]). However the critical and super-critical cases, corresponding respectively to $\alpha = 1$ and $\alpha < 1$, are harder to deal with. In the super-critical case $\alpha < 1$, we have until now only global results for small initial data, see for instance [2] and [10]. For critical case $\alpha = 1$, Constantin, C'ordoba and Wu showed in [5] the global existence in Sobolev space H^1 under smallness assumption of L^∞ norm of θ_0 but the uniqueness is proved for initial data in H^2 .

The main goal of this work is to establish a smoothing effect for a transport diffusion model in the critical case, that is when the initial data belong to the homogeneous critical Besov space.

Theorem 1.1

Let $f \in L^1_{loc}(\mathbb{R}_+, B^0_{2,1})$ and v be a smooth divergence free vector field of \mathbb{R}^2 such that $v \in L^1_{loc}(\mathbb{R}_+, Lip(\mathbb{R}^2))$. We consider also a smooth solution θ of the transport-diffusion equation (1.1), with $\alpha = \frac{1}{2}$. Let also $\theta_0 \in B^0_{2,1}$. Then there exists a positive constant c , and such that

$$\|\theta\|_{L^\infty_t B^0_{2,1}} \leq ce^{cV(t)} \left(\|\theta_0\|_{B^0_{2,1}} + \|f\|_{L^1_t B^0_{2,1}} \right),$$

where,

$$V(t) := \int_0^t \|\nabla v(\tau)\|_{L^\infty} d\tau.$$

The paper is organized as follows. In section 2, we give some definitions and recall some functional spaces. Section 3 is devoted to recall some well-known results, that will be need in the next section. In section 4, we prove our main result (Theorem 1.1).

2 Technical Tools

In this section, we recall some notations and some functional spaces as a Lebesgue space L^p , and Besov space and some results used in the paper.

2.1 Notation

- We denote by C any positive constant than will change from line to line and C_0 a real positive constant depending on the size of the initial data.
- For any A and B , we say that $A \lesssim B$, if there exist a constant $C > 0$ such that $A \leq CB$.



- The space C_0^∞ is the space of all continuous function.

2.2 Some functional spaces

This subsection is devoted to recall some functional spaces and some important results.

Definition 2.2.1

A continuous map $f: X \rightarrow Y$ is homeomorphism, if it is bijective and its inverse is continuous.

Definition 2.2.2

We define the flow associate to the velocity v by the following:

$$\psi(t, x) = x + \int_0^t v(\tau, \psi(\tau)) d\tau$$

Definition 2.2.3

we define the usual Lebesgue space $L^p(\mathbb{R}^d)$, $p \in [1, +\infty[$, by the space of all function f such that:

$$\|f\|_{L^p} := \left(\int_{\mathbb{R}^d} |f(x)|^p dx \right)^{\frac{1}{p}} < \infty,$$

and for $p = \infty$, we say that $f \in L^\infty$, if

$$\|f\|_{L^\infty} := \sup_x |f(x)| < \infty.$$

We need the definition of Besov space. We define the dyadic decomposition of the full space \mathbb{R}^2 and recall the Littlewood-Paley operators, see for example [1] and [3]. There exist two nonnegative radial functions $\chi \in \mathcal{D}(\mathbb{R}^2)$ and $\varphi \in \mathcal{D}(\mathbb{R}^2/\{0\})$ such that :

$$\chi(\xi) + \sum_{q \geq 0} \varphi(2^{-q}\xi) = 1, \quad \forall \xi \in \mathbb{R}^2,$$

$$\sum_{q \in \mathbb{Z}} \varphi(2^{-q}\xi) = 1, \quad \forall \xi \in \mathbb{R}^2/\{0\},$$

$$|p - q| \geq 2 \Rightarrow \text{supp } \varphi(2^{-p}\cdot) \cap \text{supp } \varphi(2^{-q}\cdot) = \emptyset,$$

$$q \geq 1 \Rightarrow \text{supp } \chi \cap \text{supp } \varphi(2^{-q}\cdot) = \emptyset.$$

Let $h = \mathcal{F}^{-1}\varphi$ and $\bar{h} = \mathcal{F}^{-1}\chi$, the frequency localization operators Δ_q and S_q are defined by :

$$\Delta_q f = \varphi(2^{-q}D)f, \quad S_q f = \chi(2^{-q}D)f$$

$$\Delta_{-1}f = S_0f, \quad \Delta_q f = 0 \quad \text{for } q \leq -2.$$

The homogeneous operators are defined as follows

$$\forall q \in \mathbb{Z}, \quad \dot{\Delta}_q f = \varphi(2^{-q}D)f.$$

We recall now the definition of Besov spaces, see [1] and [3].

Definition 2.2.4 (Besov space)

Let $s \in \mathbb{R}$ and $1 \leq p \leq \infty$. The inhomogeneous Besov space $B_{p,r}^s$ defined by:

$$B_{p,r}^s = \left\{ f \in \mathcal{S}'(\mathbb{R}^2) : \|f\|_{B_{p,r}^s} < \infty \right\},$$

where \mathcal{S} is the Schwartz space and

$$\|f\|_{B_{p,r}^s} := \left(2^{qs} \|\Delta_q f\|_{L^p}\right)_{l^r}.$$

The homogeneous norm :

$$\|f\|_{\dot{B}_{p,r}^s} := \left(2^{qs} \|\Delta_q f\|_{L^p}\right)_{l^r}.$$

Definition 2.2.5

Let $T > 0$ and $\rho \geq 1$, we denote by $L_T^\rho B_{p,r}^s$ the space of distribution f such that

$$\|f\|_{L_T^\rho B_{p,r}^s} := \left\| \left(2^{qs} \|\Delta_q f\|_{L^p}\right)_{\ell^r} \right\|_{L_T^\rho} < +\infty.$$

Besides the usual mixed space $L_T^\rho B_{p,r}^s$, we also need Chemin-Lerner space $\tilde{L}_T^\rho B_{p,r}^s$ which defined as the set of all distributions f satisfying

$$\|f\|_{\tilde{L}_T^\rho B_{p,r}^s} := \left\| \left\| 2^{qs} \|\Delta_q f\|_{L_T^\rho L^p} \right\|_{\ell^r} \right\| < +\infty.$$

Lemma 2.2.1 (Holder inequality)

If (f, g) belongs to $L^p \times L^q$ for any $(p, q, r) \in [1, \infty]^3$, and such that

$\frac{1}{r} = \frac{1}{p} + \frac{1}{q}$, then fg belongs to L^r and satisfies

$$\|fg\|_{L^r} \leq \|f\|_{L^p} \|g\|_{L^q}.$$

The following lemma is needed in the proof of our main result see [1] for a proof.

Lemma 2.2.2 (Gronwall's lemma)

Let f is a nonnegative continuous function on $[0, t]$, a is a real number and let A be a continuous function on $[0, t]$. Suppose also that:

$$f(t) \leq a + \int_0^t A(\tau)f(\tau)d\tau.$$

Then we have,

$$f(t) \leq a \exp\left(\int_0^t A(\tau)d\tau\right).$$

3 Some well known results

In this section, we recall some well-known results, that will be need in the next. Firstly, we give the Bernstein inequalities. This inequality, proved in [3] for any tempered distribution u , and the first author S. Sulaiman [13] and [14], proved the same inequality but for the bloc dyadic S_q and $\dot{\Delta}_q$.

Lemma 3.1 (Bernstein lemma)

There exists a constant $C > 0$ such that for all $q \in \mathbb{Z}$, $k \in \mathbb{N}$ and for every tempered distribution u we have,

$$\sup_{|\alpha|=k} \|\partial^\alpha S_q u\|_{L^b} \leq C^k 2^{q(k+2(\frac{1}{a}-\frac{1}{b}))} \|S_q u\|_{L^a}, \quad b \geq a \geq 1 \quad (2.1)$$

$$C^{-k} 2^{qk} \|\dot{\Delta}_q u\|_{L^a} \leq \sup_{|\alpha|=k} \|\partial^\alpha \dot{\Delta}_q u\|_{L^a} \leq C^k 2^{qk} \|\dot{\Delta}_q u\|_{L^a} \quad (2.2)$$

The following lemma is useful to our result, see [9], for a proof.

Lemma 3.2

Let f be any function in Schwartz class and ψ a diffeomorphism preserving Lebesgue measure, then for all $p \in [1, +\infty]$, and for all $j, q \in \mathbb{Z}$, we have

$$\|\dot{\Delta}_j(\dot{\Delta}_q f \circ \psi)\|_{L^p} \leq C 2^{|j-q|} \|\nabla \psi^{\alpha(j,q)}\|_{L^\infty} \|\dot{\Delta}_q f\|_{L^p},$$

with

$$\alpha(j, q) = \text{sign}(j - q).$$

The following result is proved in [3] and [9].

Lemma 3.3

Let u any smooth function in the Besov space $B_{2,1}^0$, and v be a divergence free vector field of \mathbb{R}^2 . Then we have

$$\sum_{q \in \mathbb{Z}} \|\Delta_q [v \cdot \nabla] u\|_{L^2} \lesssim \|\nabla v\|_{L^\infty} \|u\|_{B_{2,1}^0}.$$

The proof of the following result can be found in [9].

Proposition 3.1

If $f \in \dot{B}_{2,1}^\alpha$ such that $\alpha \in [0,1[$, and let ψ be a Lipschitz measure-preserving homeomorphism on \mathbb{R}^d . Then there exists a positive constant C_α , depend only on α , and such that

$$\| |D|^\alpha (f \circ \psi) - (|D|^\alpha f) \circ \psi \|_{L^2} \leq C e^{V(t)} V(t) 2^{\alpha q} \|f\|_{L^2},$$

with

$$V(t) := \|\nabla v\|_{L_t^1 L^\infty}.$$

The following result describes the action of the semi group operator $e^{-t|D|^\alpha}$ on distribution, whose Fourier transform is supported in a ring, see [1], [8] and [9].

Proposition 3.2

Let $p \in [1, +\infty]$, and $t, \alpha \in \mathbb{R}_+$. Then there exists a positive constant c such that for any $q \in \mathbb{N}$,

$$\|e^{-t|D|^\alpha} \Delta_q v\|_{L^p} \leq c e^{-Ct2^{q\alpha}} \|\Delta_q v\|_{L^p},$$

where the constants c and C , depend only on the dimension d .

The following can be found in [4] and [7].

Proposition 3.3

Let v be a smooth divergence free vector field. Let also f be a smooth function and θ is a smooth solution of (1.1). Then for every $p \in [1, \infty]$ we have

$$\|\theta(t)\|_{L^p} \leq \|\theta_0\|_{L^p} + \int_0^t \|f(\tau)\|_{L^p} d\tau.$$

4 Proof of our main result

We intend now to prove our main result. For this, we use a new approach based on Lagrangian coordinates combined with paradifferential calculus. The idea of the proof will be done in the spirit of [7] and [12]. First, we prove the smoothing effects for a small interval of time depending of vector v , but it depends not on the initial data.

In the second step, we proceed to division in time thereby extending the estimate at any time arbitrary chosen positive.

4.1 Local estimates

We localize in frequency the evolution equation, and rewriting the equation in Lagrangian coordinates.

Let $q \in \mathbb{N}$, then the Fourier localized function $\Delta_q \theta$ satisfies

$$\Delta_q(\partial_t \theta) + \Delta_q(v \cdot \nabla \theta) + \Delta_q(|D|^{\frac{1}{2}} + I)\theta = \Delta_q f$$

Using now the notation $[\Delta_q, v \cdot \nabla]\theta = \Delta_q(v \cdot \nabla \theta) - v \cdot \nabla \Delta_q \theta$, We get

$$\Delta_q(v \cdot \nabla \theta) = [\Delta_q, v \cdot \nabla]\theta + v \cdot \nabla \Delta_q \theta$$

This gives that

$$\partial_t \Delta_q \theta + v \cdot \nabla \Delta_q \theta + [\Delta_q, v \cdot \nabla]\theta + (|D|^{\frac{1}{2}} + I)\Delta_q \theta = \Delta_q f.$$

Therefore

$$\partial_t \Delta_q \theta + v \cdot \nabla \Delta_q \theta + (|D|^{\frac{1}{2}} + I)\Delta_q \theta = \Delta_q f - [\Delta_q, v \cdot \nabla]\theta := F_q$$

From Proposition 3.3, we have

$$\|\Delta_q \theta(t)\|_{L^2} + \|\Delta_q \theta(t)\|_{L^2}^2 \leq \|\Delta_q \theta_0\|_{L^2} + \int_0^t \|F_q(\tau)\|_{L^2} d\tau.$$

Since we have

$$\|\Delta_q \theta(t)\|_{L^2} \leq \|\Delta_q \theta(t)\|_{L^2} + \|\Delta_q \theta(t)\|_{L^2}^2 \leq \|\Delta_q \theta_0\|_{L^2} + \int_0^t \|F_q(\tau)\|_{L^2} d\tau.$$

Therefore

$$\|\Delta_q \theta(t)\|_{L^2} \leq \|\Delta_q \theta_0\|_{L^2} + \int_0^t \|F_q(\tau)\|_{L^2} d\tau.$$

Summing over q , yields

$$\sum_q \|\Delta_q \theta(t)\|_{L^2} \leq \sum_q \|\Delta_q \theta_0\|_{L^2} + \int_0^t \sum_q \|F_q(\tau)\|_{L^2} d\tau.$$

This gives that,

$$\|\theta\|_{B_{2,1}^0} \leq \|\theta_0\|_{B_{2,1}^0} + \|f\|_{L_t^1 B_{2,1}^0} + C \int_0^t \sum_q \|[\Delta_q, v \cdot \nabla] \theta(\tau)\|_{L^2} d\tau$$

Therefore, by using Lemma 3.3, we get

$$\|\theta\|_{\tilde{L}_t^\infty B_{2,1}^0} \leq \|\theta_0\|_{B_{2,1}^0} + \|f\|_{L_t^1 B_{2,1}^0} + C \int_0^t \|\nabla v(\tau)\|_{L^\infty} \|\theta\|_{\tilde{L}_\tau^\infty B_{2,1}^0} d\tau.$$

Using Lemma 2.2.2, to obtain

$$\|\theta\|_{\tilde{L}_t^\infty B_{2,1}^0} \leq C \left(\|\theta_0\|_{B_{2,1}^0} + \|f\|_{L_t^1 B_{2,1}^0} \right) e^{C \int_0^t \|\nabla v(\tau)\|_{L^\infty} d\tau}.$$

Let us now introduce the flow ψ_q of the regularized velocity v ,

$$\psi_q(t, x) = x + \int_0^t v(\tau, \psi_q(\tau, x)) d\tau.$$

We set

$$\bar{\theta}_q(t, x) = \Delta_q \theta(t, \psi_q(t, x)) \quad \text{and} \quad \bar{F}_q(t, x) = F_q(t, \psi_q(t, x)).$$

Then we have,

$$\begin{aligned} \partial_t \bar{\theta}_q + \left(|D|^{\frac{1}{2}} + I \right) \bar{\theta}_q &= \bar{F}_q + \left(|D|^{\frac{1}{2}} + I \right) (\Delta_q \theta \circ \psi_q) \\ &- \left(\left(|D|^{\frac{1}{2}} + I \right) \Delta_q \theta \right) \circ \psi_q := \bar{F}_q^1 \end{aligned} \quad (4.1)$$

Since the flow preserves Lebesgue measure, then we obtain

$$\|\bar{F}_q\|_{L^2} \leq \|\Delta_q f\|_{L^2} + \|[\Delta_q, v \cdot \nabla] \theta\|_{L^2} \quad (4.2)$$

Using now Proposition 3.1, we find that for $q \in \mathbb{Z}$

$$\left\| |D|^{\frac{1}{2}} (f \circ \psi) - \left(|D|^{\frac{1}{2}} f \right) \circ \psi \right\|_{L^2} \leq C e^{C V(t)} V(t) 2^{\frac{q}{2}} \|f\|_{L^2}, \quad V(t) := \|\nabla v\|_{L_t^1 L^\infty}.$$

This gives that,

$$\begin{aligned} & \left\| \left(|D|^{\frac{1}{2}} + I \right) (\Delta_q \theta \circ \psi_q) - \left(\left(|D|^{\frac{1}{2}} + I \right) \Delta_q \theta \right) \circ \psi_q \right\|_{L^2} \\ & \leq C e^{CV(t)} V(t) 2^{\frac{q}{2}} \|\Delta_q \theta\|_{L^2} \quad (4.3) \end{aligned}$$

Combining (4.2) and (4.3), we obtain

$$\begin{aligned} \|\bar{F}_q^1\|_{L^2} & \leq \|\Delta_q f\|_{L^2} + \|[\Delta_q, v \cdot \nabla] \theta\|_{L^2} + C e^{CV(t)} V(t) 2^{\frac{q}{2}} \|\Delta_q \theta\|_{L^2} \\ & \leq \|\Delta_q f\|_{L^2} + \|[\Delta_q, v \cdot \nabla] \theta\|_{L^2} + C e^{CV(t)} 2^{\frac{q}{2}} \|\Delta_q \theta\|_{L^2}. \end{aligned}$$

Now, we will again localize in frequency the equation (4.1) through the operator Δ_j , $j \in \mathbb{Z}$

$$\partial_t \Delta_j \bar{\theta}_q + \left(|D|^{\frac{1}{2}} + I \right) \Delta_j \bar{\theta}_q = \Delta_j \bar{F}_q^1,$$

where

$$\Delta_j \bar{F}_q^1 := \Delta_j \bar{F}_q + \Delta_j \left(\left(|D|^{\frac{1}{2}} + I \right) (\Delta_q \theta \circ \psi_q) - \left(\left(|D|^{\frac{1}{2}} + I \right) \Delta_q \theta \right) \circ \psi_q \right).$$

Using Duhamel formula,

$$\begin{aligned} \Delta_j \bar{\theta}_q(t, x) & = e^{-t(|D|^{\frac{1}{2}} + I)} \Delta_j \theta_q^0 + \int_0^t e^{-(t-\tau)(|D|^{\frac{1}{2}} + I)} \Delta_j \bar{F}_q(\tau) d\tau \\ & + \int_0^t e^{-(t-\tau)(|D|^{\frac{1}{2}} + I)} \Delta_j \left(\left(|D|^{\frac{1}{2}} + I \right) (\Delta_q \theta \circ \psi_q) - \left(\left(|D|^{\frac{1}{2}} + I \right) \Delta_q \theta \right) \circ \psi_q \right) d\tau \end{aligned}$$

Taking the L^2 norm, of the above equality, we get

$$\begin{aligned} \|\Delta_j \bar{\theta}_q(t)\|_{L^2} &\leq \left\| e^{-t(|D|^{\frac{1}{2}}+1)} \Delta_j \theta_q^0 \right\|_{L^2} + \int_0^t \left\| e^{-(t-\tau)(|D|^{\frac{1}{2}}+1)} \Delta_j \bar{F}_q(\tau) \right\|_{L^2} d\tau \\ &+ \int_0^t \left\| e^{-(t-\tau)(|D|^{\frac{1}{2}}+1)} \Delta_j \left((|D|^{\frac{1}{2}} + I) (\Delta_q \theta \circ \psi_q) - ((|D|^{\frac{1}{2}} + I) \Delta_q \theta) \circ \psi_q \right) \right\|_{L^2} d\tau. \end{aligned}$$

Using Holder inequality 2.2.1 and Proposition 3.2, we find

$$\begin{aligned} \|\Delta_j \bar{\theta}_q(t)\|_{L^2} &\leq C e^{-ct2^{\frac{j}{2}}} \|\Delta_j \theta_q^0\|_{L^2} + C \int_0^t e^{-c(t-\tau)2^{\frac{j}{2}}} \|\Delta_q f(\tau)\|_{L^2} d\tau \\ &+ C \int_0^t e^{-c(t-\tau)2^{\frac{j}{2}}} \|[\Delta_q, v \cdot \nabla] \theta(\tau)\|_{L^2} d\tau \\ &+ C e^{CV(t)2^{\frac{j}{2}}} \int_0^t e^{-c(t-\tau)2^{\frac{j}{2}}} \|\Delta_q \theta(\tau)\|_{L^2} d\tau \end{aligned}$$

Therefore,

$$\begin{aligned} \|\Delta_j \bar{\theta}_q\|_{L_t^\infty L^2} &\leq C \left(\|\Delta_j \theta_q^0\|_{L^2} + \|\Delta_q f\|_{L_t^1 L^2} \right) + \|[\Delta_q, v \cdot \nabla] \theta\|_{L_t^1 L^2} \\ &+ C e^{CV(t)2^{\frac{j}{2}}} \|\Delta_q \theta\|_{L_t^\infty L^2} \quad (4.4) \end{aligned}$$

Let $N \in \mathbb{N}$ be a fixed number that will be chosen later, and since the flow ψ preserves Lebesgue measure then we write

$$\begin{aligned} \|\Delta_q \theta\|_{L_t^\infty L^2} &= \|\bar{\theta}_q\|_{L_t^\infty L^2} \\ &\leq \sum_{|j-q| \geq N} \|\Delta_j \bar{\theta}_q\|_{L_t^\infty L^2} \\ &+ \sum_{|j-q| < N} \|\Delta_j \bar{\theta}_q\|_{L_t^\infty L^2} \\ &:= I + II. \end{aligned}$$

If $j - q \geq N$, it follows from (4.4) and by using Proposition 3.2 that,

$$\begin{aligned} \|\Delta_j \bar{\theta}_q\|_{L_t^\infty L^2} &\leq C 2^{-|j-q|} e^{\int_0^t \|\nabla v(\tau)\|_{L^\infty} d\tau} \|\Delta_q \theta\|_{L_t^\infty L^2} \\ &\leq C 2^{-|j-q|} e^{V(t)} \|\Delta_q \theta\|_{L_t^\infty L^2}. \end{aligned}$$

Therefore, we get

$$I \leq C 2^{-N} e^{CV(t)} \|\Delta_q \theta\|_{L_t^\infty L^2} \quad (4.5)$$

For the terme II ,

$$II = \sum_{|j-q| < N} \|\Delta_j \bar{\theta}_q\|_{L_t^\infty L^2},$$

we use (4.4), to get

$$II \leq C \|\Delta_q \theta_0\|_{L^2} + \|\Delta_q f\|_{L_t^1 L^2} + \|[\Delta_q, v \cdot \nabla] \theta\|_{L_t^1 L^2} + C e^{CV(t)} 2^{\frac{j}{2}} \|\Delta_q \theta\|_{L_t^\infty L^2} \quad (4.6)$$

This gives in view of (4.5) and (4.6), that

$$\|\Delta_q \theta\|_{L_t^\infty L^2} \leq I + II.$$

Thus

$$\begin{aligned} \|\Delta_q \theta\|_{L_t^\infty L^2} &\lesssim 2^{-N} e^{CV(t)} \|\Delta_q \theta\|_{L_t^\infty L^2} + \|\Delta_q \theta_0\|_{L^2} \\ &+ \|\Delta_q f\|_{L_t^1 L^2} + \|[\Delta_q, v \cdot \nabla] \theta\|_{L_t^1 L^2} + e^{CV(t)} 2^{\frac{j}{2}} \|\Delta_q \theta\|_{L_t^\infty L^2} \end{aligned}$$

Therefore,

$$\begin{aligned} \|\Delta_q \theta\|_{L_t^\infty L^2} &\lesssim \|\Delta_q \theta_0\|_{L^p} + \left[2^{-N} + 2^{\frac{j}{2}} \right] e^{CV(t)} \|\Delta_q \theta\|_{L_t^\infty L^2} \\ &+ \|\Delta_q f\|_{L_t^1 L^2} + \int_0^t \|[\Delta_q, v \cdot \nabla] \theta(\tau)\|_{L^2} d\tau. \end{aligned}$$

Now, we claim that there exists two constants $N \in \mathbb{N}$ and C_1 such that if $V(t) \leq C_1$, then

$$2^{-N} + 2^{\frac{j}{2}} \leq \frac{1}{2C}.$$

To show this, we take first t such that $V(t) \leq 1$, which is possible since $\lim_{t \rightarrow 0^+} V(t) = 0$.

Second, we choose N in order to get $2^{-N} + 2^{\frac{j}{2}} \leq \frac{1}{2C}$. By taking again $V(t)$ sufficiently small, then under this assumption $V(t) \leq C_1$, we obtain for $q \geq -1$,

$$\|\Delta_q \theta\|_{L_t^\infty L^2} \leq \|\Delta_q \theta_0\|_{L^2} + \|\Delta_q f\|_{L_t^1 L^2} + \int_0^t \|[\Delta_q, v \cdot \nabla] \theta(\tau)\|_{L^2} d\tau \quad (4.7)$$

summing over q , and using Lemma 3.3 and for $V(t) \leq C_1$, we find

$$\|\theta\|_{\tilde{L}_t^\infty B_{2,1}^0} \leq C \|\theta_0\|_{B_{2,1}^0} + C \|f\|_{L_t^1 B_{2,1}^0} + C \int_0^t \|\nabla v(\tau)\|_{L^\infty} \|\theta\|_{\tilde{L}_t^\infty B_{2,1}^0} d\tau.$$

Using Lemma 2.2.2, yields

$$\|\theta\|_{\tilde{L}_t^\infty B_{2,1}^0} \leq C \left(\|\theta_0\|_{B_{2,1}^0} + \|f\|_{L_t^1 B_{2,1}^0} \right) e^{c \int_0^t \|\nabla v(\tau)\|_{L^\infty} d\tau} \quad (4.8)$$

Therefore, the result is proved for small time. ■

4.1.2 Globalization

Let us now see how to extend this for an arbitrary positive time T . We take a partition $(T_i)_{i=0}^N$ of the interval, $[0, T]$ such that

$$\int_{T_i}^{T_{i+1}} \|\nabla v(\tau)\|_{L^\infty} d\tau \approx C_0, \forall i \in [0, N].$$

Reproducing the same argument of (4.8), we obtain

$$\|\theta\|_{\tilde{L}_{[T_i, T_{i+1}]}^\infty B_{2,1}^0} \leq C \|\theta(T_i)\|_{B_{2,1}^0} + \int_{T_i}^{T_{i+1}} \|f(\tau)\|_{B_{2,1}^0} d\tau.$$

Summing these estimates on $i = 1$, to $i = N$, and using triangle inequality, gives

$$\|\theta\|_{\tilde{L}_T^\infty B_{2,1}^0} \leq C \sum_{i=0}^{N-1} \|\theta(T_i)\|_{B_{2,1}^0} + C \int_0^T \|f(T)\|_{B_{2,1}^0} dT.$$

Thus

$$\|\theta\|_{\tilde{L}_T^\infty B_{2,1}^0} \leq CN \left(\|\theta_0\|_{B_{2,1}^0} + \|f\|_{L_T^1 B_{2,1}^0} \right) e^{CV(T)}.$$

It suffices to choose N such that $CN \approx V(t)$, then

$$\|\theta\|_{\tilde{L}_T^\infty B_{2,1}^0} \leq V(t) \left(\|\theta_0\|_{B_{2,1}^0} + \|f\|_{L_T^1 B_{2,1}^0} \right) e^{CV(T)}.$$

Therefore, we get

$$\|\theta\|_{\tilde{L}_T^\infty B_{2,1}^0} \leq C e^{CV(T)} \left(\|\theta_0\|_{B_{2,1}^0} + \|f\|_{L_T^1 B_{2,1}^0} \right).$$

This is the desired result, and the proof of the theorem is now complete. ■

Conclusion

We have proved a smoothing effects for a transport diffusion model in space dimension two. For this, we used the concept of the flow associated to the velocity and a new approach based on Lagrangian coordinates combined with paradifferential calculus.

References

- [1] Bahouri, H., Chemin, J.-Y. and Danchin, R. (2011). Fourier analysis and nonlinear partial differential equations. Springer-Verlage Berlin Heidelberg.
- [2] Chae, D. and Lee, J. (2004), Global well-posedness in the supercritical dissipative quasi-geostrophic equations, Asymptot. Anal. 38 no. 3-4, 339–358.
- [3] Chemin, J.-Y. (1998). Perfect incompressible fluids. Oxford University Press.
- [4] Cordoba, A. and Cordoba. (2004). A maximum principal applied to Quasi geostrophic equations. Com. Math. Phys, 249, n. 3, 511–528.



- [5] Constantin, P., Cordoba, D. and Wu, J. (2001). On the critical dissipative quasi-geostrophic equation, Indiana Univ. Math. J., 50, 97-107.
- [6] Constantin, P. and Wu, J. (1999). Behavior of solutions of 2D quasi-geostrophic equations, SIAM J. Math. Anal, 30, 937-948.
- [7] Hmidi, T. (2005). Regularite holderienne des poches de tourbillon visqueuses. J. Math. Pure Appl., 9, 84, n 11, 1455-1495.
- [8] Hmidi, T. and Keraani, S. (2007). Global solutions of the super-critical 2D Quasi geostrophic equations in Besov spaces, Advances in Mathematics 214 618-638.
- [9] Hmidi, T. and Keraani, S. On the global solutions of the super-critical 2D quasi-geostrophic equation in Besov spaces, <http://arxiv.org/abs/math/0611494v1>.
- [10] Ju, N. (2004). Existence and uniqueness of the solution to the dissipative 2D quasi-geostrophic equations in the Sobolev space. Comm. Math. Phus. 251, 365-376.
- [11] Pedlosky, J. (1987). Geophysical Fluid Dynamics, Springer-Verlag, New York.
- [12] Sulaiman, S. A. (2010). Global existence and uniqueness for a nonlinear Boussinesq system in dimension two. J. Math. Phys., 51, no.9, 093103.
- [13] Sulaiman, S. A. (2012). Etude qualitative de quelques equations d evolution nonlineaires. These en Univeriste de Rennes 1. France.
- [14] Sulaiman, S. A. (2021). On the Bernstein inequality, University Bulletin – ISSUE No. 22- Vol. (2) – June